

**NHS Highland**  
**Policy on treatment of breast thrush while breastfeeding**

**Directorate/Department**

**Warning – Document uncontrolled when printed**

<b>Policy Reference:</b> 010114	<b>Date of Issue:</b> March 2014
<b>Prepared by:</b> Mairi Dunbar, Lead Pharmacist – Paediatrics, Obstetrics and Gynaecology Karen MacKay – Infant Feeding Advisor	<b>Date of Review:</b> December 2020
<b>Lead Reviewer:</b> Karen Mackay	<b>Version:</b> 3
<b>Authorised by:</b> NMAHP Policies Group	<b>Date:</b>
<b>Distribution</b> <ul style="list-style-type: none"> <li>• Executive Directors</li> <li>• Clinical Directors</li> <li>• General Managers</li> <li>• Assistant General Managers</li> <li>• CHP Lead Nurses/Nurse Managers</li> <li>• Hospital Midwives</li> <li>• Community Midwives</li> <li>• Community and hospital pharmacists</li> <li>• Health Visitors</li> <li>• Public Health Practitioners</li> <li>• Nursery Nurses</li> <li>• Paediatric Nurses</li> <li>• All Paediatric Medical Staff</li> <li>• All GPs</li> <li>• All Hospital Medical Staff</li> <li>• All ancillary staff within NHS Highland</li> <li>• All support staff who have contact with mother and child</li> <li>• Neonatal unit/SCBU</li> </ul>	
<b>Method</b> CD Rom ✓      E-mail ✓                      Paper ✓                      Intranet ✓	

**Warning – Document uncontrolled when printed**

<b>Policy Reference:</b> 010114	<b>Date of Issue:</b> March 2014
<b>Prepared by:</b> Karen Mackay, Mairi Dunbar	<b>Date of Review:</b> December 2020
<b>Lead Reviewer:</b> Karen Mackay	<b>Version:</b> 3
<b>Authorised by:</b> NMAHP Policies Group	<b>Page</b> 1 of 6

## CONTENTS

Introduction	3
Symptoms	3
Prescribing for thrush in lactated women	5
References	6

**Warning – Document uncontrolled when printed**

<b>Policy Reference:</b> 010114	<b>Date of Issue:</b> March 2014
<b>Prepared by:</b> Karen Mackay, Mairi Dunbar	<b>Date of Review:</b> December 2020
<b>Lead Reviewer:</b> Karen Mackay	<b>Version:</b> 3
<b>Authorised by:</b> NMAHP Policies Group	<b>Page</b> 2 of 6

## INTRODUCTION

### What is Thrush? (*Candida albicans*)

Thrush is a fungal infection and can occur any time while the mother is breastfeeding. *Candida albicans* thrive in moist, warm, dark areas and the breasts are ideal to harbour this fungus. *Candida* infections most commonly occur where there has been a breakdown in the integrity of the skin or mucous membrane. The widespread use of antibiotics now in the ante and intra-partum period encourages the overgrowth of *Candida albicans*.

### Diagnosis of breast thrush

There is no real diagnostic test available, although the breastfeeding network (BFN) 2013 have suggested taking a black charcoal media swab from both mother's nipples and baby's mouth to confirm presence of either *Candida albicans* or bacterial infection.

It is vitally important that a clear and in-depth breastfeeding history is taken and a breastfeed is observed by a trained health professional. In doing so, other potential problems can be eliminated and a clear thrush diagnosis can be made. It is also important to consider that there will be cases where a mother will display symptoms and signs of thrush while the baby will be symptom free and vice versa.

## SYMPTOMS

### For the Breastfeeding Mother

- Nipple pain that begins after a period of pain free feeding.
- Pain is not improved by effective attachment.
- Cracked nipples which are difficult and slow to heal.
- Pain is often described as severe in nature – burning and stabbing.
- Pain commences during the feed – not at the start and continues after the feed has ended, sometimes for as long as an hour.
- Loss of colour to the nipple or areola.
- Sensitive nipples and areola.
- History of recent antibiotic therapy.
- History of recent episode of vaginal thrush.
- Pain will always occur in **both** breasts due to the baby transferring the infection from one to the other breast.
- Absence of a pyrexia.

### For the Baby When Breastfeeding

- Inability to sustain a latch due to oral pain.
- Frustration at breast – going on and off.
- Crying.
- Nappy rash which doesn't improve with simple treatment.
- Can have oral plaques in mouth.
- Whitish sheen to saliva and inside the lips and gums (Mohrbacher and Stock 2003).
- Breast refusal.
- Colic symptoms: infants may be windy, fretful and find it hard to settle (The breastfeeding Network 2003a).
- Poor weight gain.
- A baby with oral coating to posterior part of mouth only with no other symptoms should be assessed for tongue tie

Warning – Document uncontrolled when printed

Policy Reference: 010114	Date of Issue: March 2014
Prepared by: Karen Mackay, Mairi Dunbar	Date of Review: December 2020
Lead Reviewer: Karen Mackay	Version: 3
Authorised by: NMAHP Policies Group	Page 3 of 6

## **Note – infants may have no symptoms**

If pain is isolated to one breast then this is not symptomatic of thrush and assessment of a full breastfeed is vital. A bacteriology swab is recommended for a unilateral breast pain where alteration of positioning and attachment has not helped to rule out a staph aureus infection which may need treatment with antibiotics

It is vital for a health professional trained in breastfeeding management to observe a full breastfeed prior to diagnosing thrush and/or commencing thrush treatment. This will usually be done by either the local midwife or health visitor. The reasoning behind this is to rule out any other causes for the mother's symptoms. When referring a mother to the G.P, the health professional is encouraged to have observed a full breastfeed prior to referring to G.P.

Effective positioning and attachment is crucial to ensure adequate milk production, milk transfer and breast health. This should be the primary concern when assessing any breastfeeding problem. Once effective positioning and attachment is confirmed then other causes of nipple pain can be considered.

### **Other causes of Nipple Pain**

- Eczema
- Tongue tie in the baby
- Reynaud's syndrome
- Milk bleb or white spot
- Bacterial infection which will may appear as a sloughy discharge

### **Self help to treat thrush**

- Pain killers are encouraged to relieve the pain experienced from thrush – over the counter analgesics such as paracetamol and ibuprofen are ideal.
- Continue to breastfeed as thrush will not harm your baby.
- Personal hygiene is important to stop cross –infection. Good hand washing from all the family is required.
- If a dummy, nipple shield, teat, bottle or plastic toys are used by the baby ensure that these are carefully washed and sterilised while the infection lasts – if the mother has other children make sure that their toys that come in to contact with the baby are carefully washed too to save cross contamination.
- Any milk which has been expressed and stored in the freezer during the time when you have had thrush should be discarded.
- Some mothers find reducing the amount of sugar and yeast in their diet helps but this has not been researched.

#### **Warning – Document uncontrolled when printed**

<b>Policy Reference:</b> 010114	<b>Date of Issue:</b> March 2014
<b>Prepared by:</b> Karen Mackay, Mairi Dunbar	<b>Date of Review:</b> December 2020
<b>Lead Reviewer:</b> Karen Mackay	<b>Version:</b> 3
<b>Authorised by:</b> NMAHP Policies Group	<b>Page</b> 4 of 6

## PRESCRIBING FOR THRUSH IN LACTATING WOMEN

### BOTH MUM AND BABY MUST BE TREATED TOGETHER

#### Surface thrush

- Miconazole 2% Cream for surface thrush. Best practice is to apply a small amount more frequently after every feed rather than the licensed twice daily treatment. Any cream which can be seen should be wiped off gently prior to the next feed to reduce further nipple damage. If no cream is visible then no need to wipe off<sup>1,2</sup>. Use for 10 days after the symptoms have subsided or more specialist help is obtained. The baby should also be treated as per below.

#### Ductal thrush

- In addition to miconazole 2% cream; Fluconazole orally 150mg as a loading dose then 100mg daily for at least 10 days<sup>2,3,4,5,10</sup>.
- If pain has not improved in 2 to 3 days, increase dose to 200mg daily.
- If symptoms are very severe or prolonged, give doses up to 400mg as a loading dose followed by 200mg daily for 10 days<sup>2</sup>.

#### Prescribing for oral thrush in babies

- **First line treatment:** Neonates and children – Nystatin 100,000 units/ml oral suspension, 1ml four times daily after feeds or food<sup>6,10</sup>.
- **Second line treatment:** - Miconazole oral gel smeared around the inside of the mouth neonate 1 ml four times a day after feeds,; 1 month to 1 year – 1.25ml four times a day after feeds or food; from 2 years – 2.5ml four times a day(**unlicensed under 4 months**)<sup>1,6</sup>. Apply to all areas of the mouth, cheeks, roof of mouth and the tongue with a clean fingertip. Ensure that the gel does not obstruct the mouth. DO NOT apply the gel to the back of the throat, and the full dose may need to be divided into smaller portions<sup>7</sup>. Continue for at least 7 days after lesions have healed or symptoms have cleared. Nystatin suspension does not adhere so well as Miconazole oral gel<sup>1,2</sup>.
- Nurse prescribers can prescribe these treatments.

**NOTE: the Summary of Product Characteristics (SPC) for Miconazole (Daktarin) oral gel does not recommend the use of oral gel in any patient less than four months old (and cautions against use in any pre-term infant less than 6 months old)<sup>7</sup>. This could be because of a case report in 2004 when a 17 day old child in the Netherlands, born at 36 weeks, choked on some of the viscous gel and gel was then removed from the back of the throat. There were another 9 cases reported in the Netherlands, 7 of which occurred in babies less than 2 months old.**

#### Use of Miconazole and Fluconazole in breastfeeding

- Miconazole Cream is not absorbed through the skin to a great extent, and topical use during breastfeeding requires no caution in use<sup>3,4</sup>.
- Fluconazole passes into breast milk to reach concentrations lower than those in plasma<sup>8</sup>. It is licensed

Warning – Document uncontrolled when printed

Policy Reference: 010114	Date of Issue: March 2014
Prepared by: Karen Mackay, Mairi Dunbar	Date of Review: December 2020
Lead Reviewer: Karen Mackay	Version: 3
Authorised by: NMAHP Policies Group	Page 5 of 6

to be used in breastfeeding after a single use of a standard dose 200 mg or less<sup>8</sup>. It is not licensed in breastfeeding after repeated use or after high doses<sup>8</sup>. An infant will receive between 16 to 22% of the weight adjusted maternal dose which is equivalent to about 0.34 to 0.5mg/kg/day for the infant<sup>3,4</sup>. This is well below therapeutic doses of e.g. 3 to 12mg/kg every 72 hours, 48 hours, or daily, depending on the age of the child<sup>6</sup>. Preterm neonates weighing less than 1000g have received fluconazole prophylactically and have demonstrated successful outcomes<sup>9</sup>. The World Health Organisation recognises fluconazole as compatible with breastfeeding, although there is potential for accumulation in a neonate < 6 weeks. The fluconazole dose should be taken after the last breast feed if possible.

## REFERENCES

1. <http://intranet.nhsh.scot.nhs.uk/Org/CorpServ/PublicHealth/InfantFeeding/Documents/Breast%20Feeding%20Problems%201105%20no%202.ppt>
2. <https://breastfeedingnetwork.org.uk/wp-content/dibm/thrush%20detailed%20information%20and%20breastfeeding.pdf>
3. Drugs during Pregnancy and Lactation. Schaefer, Peters and Miller. 3<sup>rd</sup> edition.
4. Medications and Mothers' Milk 2014. Thomas W. Hale.
5. Neonatal Formulary. Drug use in pregnancy and the first year of life. 7<sup>th</sup> Edition.
6. BNF for Children. [www.bnfc.org](http://www.bnfc.org)
7. <http://www.medicines.org.uk/emc/medicine/7301>
8. <http://www.medicines.org.uk/emc/medicine/25676>
9. Fluconazole prophylaxis against fungal colonisation and infection in preterm infants. Kaufman D. et al. N Eng J Med 2001;345(23): 1660-6.
10. Highland Formulary 7th edition

**Warning – Document uncontrolled when printed**

<b>Policy Reference:</b> 010114	<b>Date of Issue:</b> March 2014
<b>Prepared by:</b> Karen Mackay, Mairi Dunbar	<b>Date of Review:</b> December 2020
<b>Lead Reviewer:</b> Karen Mackay	<b>Version:</b> 3
<b>Authorised by:</b> NMAHP Policies Group	<b>Page</b> 6 of 6